

To appear in ApJ (Letters)

Compact Centimeter and Millimeter Sources in NGC 6334 I(N): OB Stars in the Making?

Luis F. Rodríguez and Luis A. Zapata¹

Centro de Radioastronomía y Astrofísica, UNAM, Morelia 58089, México

`l.rodriguez, l.zapata@astrosmo.unam.mx`

and

Paul T. P. Ho¹

Academia Sinica Institute of Astronomy and Astrophysics, Taipei 106, Taiwan

`pho@asiaa.sinica.edu.tw`

ABSTRACT

We present sensitive, high angular resolution 1.3 cm and 7 mm observations of the massive core NGC 6334 I(N), a region known to be undergoing massive star formation. At 1.3 cm we detect three sources, of which two had been previously detected at centimeter or millimeter wavelengths. At 7 mm we detect four sources. We suggest that three of these sources are subcomponents of the millimeter source SMA 1, that at these wavelengths is the dominant source in the region. The fourth 7 mm source appears to be associated with the relatively isolated source SMA 6. In all four 7 mm sources, the continuum emission is arising from structures of dimensions in the order of 1000 AU for which we estimate masses of order a few solar masses. We interpret these 7 mm sources as massive circumstellar disks that, however, surround stars or compact small stellar groups that at present have masses comparable to those of the disks but that may be accreting on their way to become massive stars.

Subject headings: stars: formation — ISM: individual (NGC 6334 I(N))

¹Harvard-Smithsonian Center for Astrophysics, and Submillimeter Array, 60 Garden Street, Cambridge, MA 02138, USA

1. Introduction

Located at the NE of the molecular cloud/H II region complex NGC 6334, the massive cores NGC 6334 I and NGC 6334 I(N) are separated by only $90''$ and are considered “twin” objects in some aspects but also very different in other characteristics (Beuther et al. 2005). Both cores have similar dimensions: $\sim 10''$ (Sandell 2000) or 0.08 pc at a distance of 1.7 kpc (Neckel 1978). They also have similar masses in gas ($\sim 200 M_{\odot}$ for NGC 6334 I and $\sim 400 M_{\odot}$ for NGC 6334 I(N); Sandell 2000), but their bolometric luminosities are very different ($\sim 2.6 \times 10^5 L_{\odot}$ for NGC 6334 I and $\sim 1.9 \times 10^3 L_{\odot}$ for NGC 6334 I(N); Sandell 2000).

These differences have been attributed to NGC 6334 I(N) being less evolved than NGC 6334 I (Thorwirth et al. 2003; Hunter et al. 2006). With time, the forming massive stars in NGC 6334 I(N) will reach their final masses and shine with much larger luminosity than now. There is recent observational evidence that the formation of massive stars takes place in a similar way to that of solar type objects, namely by accretion via a circumstellar disk: a handful of cases where a disk could be present around a forming massive star have been presented in the literature (e. g. Shepherd & Kurtz 1999; Patel et al. 2005; Schreyer et al. 2006). These disks are larger than those found in forming solar stars, and have dimensions of order 1000 AU and masses of a few solar masses. In this paper we present 1.3 cm and 7 mm observations of the core NGC 6334 I(N), trying to study the environment of the massive stars that are suspected to be forming there.

2. Observations

The 1.3 cm observations were made in the BnA configuration of the VLA of the NRAO¹, during 2005 January 23. The central frequency observed was 22.46 GHz and we integrated on-source for a total of 120 minutes. The absolute amplitude calibrator was 1331+305 (with an adopted flux density of 2.52 Jy) and the phase calibrator was 1720–358 (with a bootstrapped flux density of 0.652 ± 0.005 Jy). The phase noise rms was about 25° , indicating good weather conditions. The phase center of these observations was at $\alpha(2000) = 17^h 20^m 54^s.9$; $\delta(2000) = -35^{\circ} 44' 54''$. The 7 mm observations were made in the CnB configuration of the VLA during 2005 June 30. The central frequency observed was 43.34 GHz and we integrated on-source for a total of 100 minutes. The absolute amplitude calibrator was 1331+305 (with an adopted flux density of 1.45 Jy) and the phase calibrator

¹The National Radio Astronomy Observatory is operated by Associated Universities Inc. under cooperative agreement with the National Science Foundation.

was 1720–358 (with a bootstrapped flux density of 1.02 ± 0.01 Jy). The phase noise rms was about 10° , indicating excellent weather conditions. The phase center of these observations was at $\alpha(2000) = 17^h 20^m 54^s.9$; $\delta(2000) = -35^\circ 45' 06''$.

The data were acquired and reduced using the recommended VLA procedures for high frequency data, including the fast-switching mode with a cycle of 120 seconds. The effective bandwidth of the observations was 100 MHz. We made images with natural weighting to obtain the highest signal-to-noise possible.

3. Analysis

3.1. 1.3 cm Data

The rms noise of the 1.3 cm image is $67 \mu\text{Jy}$. At this wavelength we detected three sources, whose positions and total flux densities are given in Table 1. All three sources appear unresolved ($\leq 0''.2 - 0''.3$). The first source in Table 1 does not have a counterpart at other wavelengths and we refer to it as VLA-K 1 (since it was detected at 1.3 cm, the so-called K band). The other two sources are associated, respectively, with the sources SMA 4 and SMA 1 reported by Hunter et al. (2006). In Figure 1 we show a contour image of these last two sources as well as the positions of other sources in the region.

3.2. 7 mm Data

The rms noise of the 7 mm image is $320 \mu\text{Jy}$. At this wavelength we detected four sources, whose positions and total flux densities are given in Table 2. The first source in Table 2 is associated with SMA 6, a source reported by Hunter et al. (2006). In Figure 2 we show a contour image of this 7 mm source. The other three 7 mm sources are very close among them, forming a line that extends by $\sim 4''$ (see Figure 3). We interpret these three sources as the subcomponents of the 1.3 mm source SMA 1, the dominant millimeter object in the region (Hunter et al. 2006). SMA 1 extends by about the same extent and position angle as the three 7 mm sources. We then refer to the three 7 mm sources as SMA 1a, 1b, and 1c (see Table 2).

3.3. The Nature of the Continuum Emission

Of the three sources detected at 1.3 cm, only one (SMA 4) has an associated 3.6 cm counterpart, an unresolved source with flux density of 0.34 ± 0.06 mJy detected by Carral et al. (2002). The other two sources detected at 1.3 cm do not have a 3.6 cm counterpart at the $4\text{-}\sigma$ level of ~ 0.24 mJy. The spectral index between 3.6 cm and 1.3 cm is then 0.4 ± 0.3 ($S_\nu \propto \nu^{0.4 \pm 0.3}$), consistent with free-free emission from a thermal jet (Reynolds 1986), although an optically-thin HII region cannot be ruled out. This spectral index determination is reliable since the 3.6 and 1.3 cm observations have similar angular resolution ($\sim 0''.2 - 0''.3$). The other two 1.3 cm sources do not have a 3.6 cm counterpart and lower limits of ~ 0.9 are estimated for their spectral indices. Again, this spectral index is consistent with the values expected for free-free emission from a thermal jet. We suggest that the compact 1.3 cm emission detected by us is probably tracing ionized jets.

Of the four sources detected at 7 mm, only one (SMA 1b) has an associated 1.3 cm counterpart, an unresolved source with flux density of 0.57 ± 0.12 mJy detected by us (see Tables 1 and 2 and Figure 1). The spectral index of this source between 1.3 cm and 7 mm is large, 4.5 ± 0.4 , suggesting that the 7 mm emission cannot be attributed to optically-thick free-free emission and is most probably arising from dust. A similar conclusion is reached for the other three 7 mm sources that are not detected at 1.3 cm, but show a flux density of several mJy at 7 mm. The non detection of these three sources at 1.3 cm, indicates that their spectral index between 7 mm and 1.3 cm is ≥ 3.3 . We then suggest that, in contrast to the 1.3 cm emission that appears to be tracing free-free emission, the 7 mm emission is tracing dust. The 1.3 cm data has better angular resolution ($\sim 0''.3$) than the 7 mm data ($\sim 0''.6$). However, we have tapered the 1.3 cm data to produce images of similar angular resolution to those at 7 mm and our results are not changed.

We finally compare our 7 mm data with the 1.3 mm data of Hunter et al. (2006). This comparison is not direct since our angular resolution is of order $0''.6$ while that of Hunter et al. (2006) is of order $1''.6$. In particular, as noted before, the source SMA 1 of Hunter et al. (2006) that is the dominant 1.3 mm source of the region is resolved by us into three subcomponents at 7 mm (SMA 1a, 1b, and 1c). The total flux density of SMA 1 at 1.3 mm is 2.04 Jy. In a solid angle similar to that occupied by SMA 1 at 1.3 mm, we detect a total flux density of ~ 40 mJy at 7 mm (of which 31 mJy can be attributed to the sources SMA 1a, 1b, and 1c). A spectral index of 2.4 ± 0.2 is derived between 7 and 1.3 mm. A similar comparison can be made for the source SMA 6, for which we estimate a total flux density of ~ 9 mJy for a solid angle similar to that occupied by SMA 6 at 1.3 mm. The same spectral index, 2.4 ± 0.2 , as derived for SMA 1 is obtained for SMA 6 between 7 and 1.3 mm. We then suggest that at 7 and 1.3 mm we are observing optically thin dust emission with a dust

mass opacity coefficient that goes with frequency as $\kappa_\nu \propto \nu^{0.4}$. With this information we can estimate the masses of the 7 mm sources. Following Hunter et al. (2006) we adopt a value of $\kappa_{1.3mm} = 1.5 \text{ cm}^2 \text{ g}^{-1}$ (the average of the values of $1.0 \text{ cm}^2 \text{ g}^{-1}$, valid for grains with thick dust mantles, and $2.0 \text{ cm}^2 \text{ g}^{-1}$, valid for grains without mantles). This implies $\kappa_{7mm} = 0.78 \text{ cm}^2 \text{ g}^{-1}$. Assuming optically thin, isothermal dust emission and a gas-to-dust ratio of 100 (Sodroski et al. 1997), the total mass of the 7 mm sources is given by:

$$\left[\frac{M}{M_\odot} \right] = 0.11 \left[\frac{S_\nu}{mJy} \right] \left[\frac{T}{100 \text{ K}} \right]^{-1} \left[\frac{\nu}{43 \text{ GHz}} \right]^{-2.4} \left[\frac{D}{kpc} \right]^2,$$

where T is the dust temperature and D is the distance to the source. Assuming $T = 100 \text{ K}$ for SMA 1 and $T = 33 \text{ K}$ for SMA 6 (Hunter et al. 2006), we derive masses on the order of a few solar masses for the four 7 mm sources (see Table 2). These dust temperatures (Hunter et al. 2006) have been estimated from molecular observations and have uncertainties of a factor of two, that reflect in the mass determination.

We finally discuss the different spectral indices for the emission of the sources determined by us for the 1.3 mm to 7 mm range (that gives a value of ~ 2.4), and for the 7 mm to 1.3 cm range (that gives a value of ≥ 3.3). As noted by Wilner et al. (2005), these changes in spectral index most probably denote grain growth to centimeter-size particles. The significant presence of centimeter-size dust grains flattens the spectral index for wavelengths shorter than the characteristic size of the grains.

3.4. Massive Circumstellar Disks?

The dimensions ($\sim 1000 \text{ AU}$) and masses (a few $M_\odot$) of the four sources detected at 7 mm are suggestive of massive circumstellar disks. Furthermore, the two 7 mm sources with deconvolved dimensions (SMA 1a and SMA 1b; see Table 2) have orientations consistent with hosting the exciting sources of two outflows reported in the region. Megeath et al. (1999) detected an SiO outflow whose axis is oriented at position angle of $\sim 135^\circ$ and could be attributed to the source associated with SMA 1a, whose major axis is oriented at a position angle of $\sim 42^\circ$, nearly perpendicular to the outflow. Hunter et al. (2006) propose that an elongated infrared nebulosity could be a jet or a reddened reflection nebulosity from an outflow cavity and propose that it is excited by SMA 4. This nebulosity extends approximately to the west of the core of NGC 6334 I(N), with position angle of $\sim 240^\circ$ and could alternatively be excited by SMA 1b, since this source (as SMA 4) also lies along the axis of the nebulosity.

There is, however, a problem with the interpretation of these structures as massive circumstellar disks. In the cases of forming objects where the masses of the disk and its central star have been estimated, they usually have a ratio of order $M_{star}/M_{disk} \simeq 10$ (Rodríguez et al. 1998; Guilloteau & Dutrey 1998; Schreyer et al. 2006). If we apply this mass ratio to the 7 mm sources in NGC 6334 I(N), stars with masses of tens of solar masses would be required at the center of the disks. This is clearly not the case since the modest bolometric luminosity of $\sim 1.9 \times 10^3 L_{\odot}$ (Sandell 2000) would then be exceeded by one to two orders of magnitude.

However, the ratio of $M_{star}/M_{disk} \simeq 10$ is not an established rule. For forming objects, surely the M_{star}/M_{disk} ratio will depend on the stage of evolution, with smaller values at the initial stages. The theoretical models of Yorke, Bodenheimer, & Laughlin (1995) predict values of $M_{star}/M_{disk} \simeq 4$ for forming intermediate-mass stars. It is known from theoretical two-dimensional calculations that forming stars with $M_{star}/M_{disk} \leq 3$ will face gravitational instabilities that will induce spiral waves (Laughlin & Bodenheimer 1994). However, from their global, three-dimensional smoothed particle hydrodynamics simulations Lodato & Rice (2005) conclude that these massive self-gravitating disks can survive for relatively long periods. The problem of disk stability in massive stars has been recently reviewed by Cesaroni et al. (2006).

From the observational point of view, in some sources the possibility that M_{star}/M_{disk} is as low as ~ 2 is consistent with the data (Patel et al. 2005). Furthermore, in the case of G192.16-3.82, the protostar and disk have comparable masses of about $8 M_{\odot}$ (Shepherd, Claussen, & Kurtz 2001). With the finding of additional young objects that may have comparable masses in the disk and the protostar, it is important to review the issue of the stability of these structures. It is also possible that these massive disks are forming a compact stellar group, as proposed by Sollins & Ho (2005) for the $100 M_{\odot}$ disk in G10.6-0.4, in which case part of the mass now observed in them will end in the smaller members of the group.

In summary, we believe that the observations discussed here are consistent with the presence in NGC 6334 I(N) of several forming massive stars that at present have reached only a fraction of their final masses and are surrounded by disks with masses comparable to those of the stars. Whether these massive structures that we identify as disks are forming a single star or a compact stellar group cannot be disentangled with the present observations.

LFR and LAZ acknowledge the support of CONACyT, México and DGAPA, UNAM.

Facilities: VLA

REFERENCES

- Beuther, H., Thonwirth, S., Zhang, Q., Hunter, T. R., Megeath, S. T., Walsh, A. J., & Menten, K. M. 2005, *ApJ*, 627, 834
- Carral, P., Kurtz, S. E., Rodríguez, L. F., Menten, K., Cantó, J., & Arceo, R. 2002, *AJ*, 123, 2574
- Cesaroni, R., Galli, D., Lodato, G., Walmsley, C. M. & Zhang, Q. 2006, in *Protostars and Planets V*, ed. B. Reipurth, D. Jewitt, & K. Keili (Tucson: Univ. Arizona Press), in press
- Guilloteau, S. & Dutrey, A. 1998, *A&A*, 339, 467
- Hunter, T. R., Brogan, C. L., Megeath, S. T., Menten, K. M., Beuther, H., & Thorwirth, S. 2006, submitted to *ApJ*.
- Laughlin, G. & Bodenheimer, P. 1994, *ApJ*, 436, 335
- Lodato, G. & Rice, W. K. M. 2005, *MNRAS*, 358, 1489
- Megeath, S. T. & Tieftrunk, A. R. 1999, *ApJ*, 526, L113
- Neckel, T. 1978, *A&A*, 69, 51
- Patel, N. A., Curiel, S., Sridharan, T. K., Zhang, Q., Hunter, T. R., Ho, Paul T. P., Torrelles, J. M., Moran, J. M., Gómez, J. F., & Anglada, G. 2005, *Nature*, 437, 109
- Reynolds, S. P. 1986, *ApJ*, 304, 713
- Rodríguez, L. F., D'Alessio, P., Wilner, D. J., Ho, P. T. P., Torrelles, J. M., Curiel, S., Gómez, Y., Lizano, S., Pedlar, A., Cantó, J., & Raga, A. C. 1998, *Nature*, 395, 355
- Sandell, G. 2000, *A&A*, 358, 242
- Schreyer, S., Semenov, D., Henning, Th., & Forbrich, J. 2006, *ApJ*, 637, L129
- Shepherd, D. S. & Kurtz, S. E. 1999, *ApJ*, 523, 690
- Shepherd, D. S., Claussen, M. J., & Kurtz, S. E. 2001, *Science*, 292, 1513
- Sodroski, T. J., Odegard, N., Arendt, R. G., Dwek, E., Weiland, J. L., Hauser, M. G., & Kelsall, T. 1997, *ApJ*, 480, 173
- Sollins, P. K. & Ho, P. T. P. 2005, *ApJ*, 630, 987

- Thorwirth, S., Winnewisser, G., Megeath, S. T., & Tieftrunk, A. R. 2003, in ASP Conf. Ser. 287, *Galactic Star Formation Across the Stellar Mass Spectrum*, ed. J. M. De Buizer & N. S. van der Blik (San Francisco: ASP), 257
- Wilner, D. J., D’Alessio, P., Calvet, N., Claussen, M. J., & Hartmann, L. 2005, *ApJ*, 626, L109
- Yorke, H. W., Bodenheimer, P., & Laughlin, G. 1995, *ApJ*, 443, 199




Fig. 1.— Contour image of the 1.3 cm emission of the core of NGC 6334 I(N). Two of the three sources detected at this wavelength, SMA 4 and SMA 1b, are included in this region (see Table 1 for the positions and flux densities of these three sources). Contours are -4, 4, 5, and 6 times $67 \mu\text{Jy}$, the rms noise of the image. The half power contour of the synthesized beam ($0''.38 \times 0''.27$ with a position angle of $+19^\circ$) is shown in the bottom left corner. The center of the empty square marks the position of one of the two 3.6 cm sources detected by Carral et al. (2002). The crosses mark the position and positional error of five of the seven 1.3 mm sources reported by Hunter et al. (2006), while the asterisks mark the position of three of the four 7 mm sources reported in this paper.




Fig. 2.— Contour image of the 7 mm emission of the source associated with the millimeter object SMA 6. Contours are -3, 3, 4, and 5 times $320 \mu\text{Jy}$, the rms noise of the image. The half power contour of the synthesized beam ($0''.69 \times 0''.44$ with a position angle of $+25^\circ$) is shown in the bottom left corner. The cross marks the position and positional error of the source SMA 6 detected by Hunter et al. (2006).




Fig. 3.— Contour image of the 7 mm emission of the core of NGC 6334 I(N). Three of the four sources detected at this wavelength, SMA 1a, SMA 1b, and SMA 1c, are included in this region (see Table 2 for the positions and flux densities of these four sources). Contours are -3, 3, 4, 5, and 6 times 320 μ Jy, the rms noise of the image. The half power contour of the synthesized beam ($0''.69 \times 0''.44$ with a position angle of $+25^\circ$) is shown in the bottom left corner. The dashed lines show the orientations of two outflows in the region that could be associated with the sources SMA 1a and SMA 1b. The center of the empty diamond marks the position of one of the three 1.3 cm sources reported in this paper (see Fig. 1 and Table 1). The crosses mark the position and positional error of four of the seven 1.3 mm sources reported by Hunter et al. (2006).

Table 1: Parameters of the 1.3 cm Sources in NGC 6334 I(N).

Source	Position ^a			Total Flux
	α (J2000)	δ (J2000)		Density ^b (mJy)
VLA-K 1	17 20 52.701	-35 45 13.52		0.59±0.09
SMA 4	17 20 54.622	-35 45 08.23		0.50±0.11
SMA 1b	17 20 55.192	-35 45 03.54		0.57±0.12

^aUnits of right ascension are hours, minutes, and seconds and units of declination are degrees, arcminutes, and arcseconds. Positional accuracy is estimated to be 0''05.

^bAll three sources appear unresolved, with angular size $\leq 0''.2 - 0''.3$.

Table 2: Parameters of the 7 mm Sources in NGC 6334 I(N).

Source	Position ^a		Total Flux		Deconvolved Angular Size ^b	Estimated Mass (<i>M</i> _⊙)
	α(J2000)	δ(J2000)	Density (mJy)			
SMA 6	17 20 54.594	-35 45 17.01	3.8±1.0		≤ 1''0	4
SMA 1a	17 20 55.170	-35 45 05.26	13.6±2.4		1''9 ± 0''5 × 1''3 ± 0''4; + 42° ± 36°	4
SMA 1b	17 20 55.199	-35 45 03.43	11.0±1.8		1''4 ± 0''4 × 1''2 ± 0''5; + 164° ± 33°	3
SMA 1c	17 20 55.260	-35 45 02.42	6.8±1.5		≤ 1''0	2

^aUnits of right ascension are hours, minutes, and seconds and units of declination are degrees, arcminutes, and arcseconds. Positional accuracy is estimated to be 0''.2.

^bMajor axis $\times$ minor axis; position angle of major axis.